

POST OPERATIVE INFORMATION

Shoulder Surgery

Normal Symptoms:

- · Discomfort in shoulder
- Mild Swelling
- Bruising

The following symptoms are **NOT** normal:

- · Prolonged vomiting
- Excessive bleeding
- · Redness around the incision
- · Fever or chills
- · Increasing swelling or numbness

If you experience any of the above symptoms you should seek medical advice. Please contact your nearest Emergency Department.

Diet:

Normal healthy diet as tolerated.

Activity:

- Walking for exercise.
- Must wear sling for the next six weeks until reviewed by Doctor.
- · No heavy lifting.

Medications:

- If you have any pain or swelling put ice on your shoulder and rest.
- You may use oral analgesia (eg Panadol) as required.

Driving:

No driving until instructed by Doctor.

Dressings / Wound care:

- · Wash and pat wounds dry carefully.
- Leave the dressing in place until seen by Doctor.

Hygiene:

· You may shower as desired.

Legal requirements following an anaesthetic or sedation:

For a 24 hour period you are NOT allowed to:

- Drive a car as directed by Doctor
- Operate machinery
- · Drink alcohol
- · Sign any legal or important documents.

Follow-up Appointment





DR L. ANDREW ASHTON

B.A., B.Med., FRACS (Orth) Orthopaedic Surgeon

E secretary@oso.com.au T 0263 601 500 309 Anson Street, Orange NSW 2800 PO Box 8136, East Orange 2800 www.orthopaedicsurgeonorange.com.au

PROVIDER NO. 204803DF **ABN** 13371125435